

N-Acetyl Aspartyl Glutamic Acid Sodium Salt

Naabak®

49 mg / mL (4.9%)

Ophthalmic Solution (Drops)

Anti-allergy

FORMULATION:

Each mL contains 49 mg of N-Acetyl Aspartyl Glutamic Acid Sodium Salt.

THERAPEUTIC INDICATIONS:

This medicinal product is recommended for the relief of ocular symptoms of allergic origin (conjunctivitis and blepharoconjunctivitis).

This product is **FOR EXTERNAL USE ONLY**

WARNING!

CASES IN WHICH THIS MEDICINAL PRODUCT SHOULD NOT BE USED (CONTRAINDICATIONS):

IN DOUBT, SEEK ADVICE FROM YOUR DOCTOR OR PHARMACIST

SPECIAL WARNINGS:

- Do not inject, do not swallow
- The absence of preservative in the instilled drops allows contact lens wearing
- Do not touch the eye with the bottle tip.

PRECAUTIONS FOR USE:

- In children under 4, DO NOT USE WITHOUT MEDICAL ADVICE.
- In case of severe conjunctival allergy, SEEK ADVICE FROM YOUR DOCTOR
- Do not exceed the prescribed or recommended dosage, do not stop your treatment without medical advice.

IN DOUBT, DO NOT HESITATE TO SEEK ADVICE FROM YOUR DOCTOR OR PHARMACIST

DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS:

IN ORDER TO AVOID POSSIBLE INTERACTION BETWEEN DIFFERENT MEDICINAL PRODUCTS, NOTIFY YOUR DOCTOR OR PHARMACIST OF ANY OTHER CONCOMITANT TREATMENTS.

PREGNANCY-BREAST-FEEDING

ALWAYS SEEK ADVICE FROM YOUR DOCTOR OR PHARMACIST PRIOR TO USING A MEDICINAL PRODUCT

HOW TO USE THIS MEDICINAL PRODUCT?

DOSAGE: 1 Drop, 2 to 6 times daily or as prescribed by the physician

APPLICATION

Topical application - OCULAR INSTILLATION

To ensure a good use of this eye drops, some precautions must be taken:

- Carefully wash your hands prior to proceed to the applications
- Avoid contact between the eye or the lids of the bottle tip. Instill one drop of the solution into the conjunctival sac while pulling the lower eyelid downwards and looking up. The appearance time of a drop could be longer than with a standard eye drops solution. Screw down the bottle immediately after instillation.

FREQUENCY AND TIME OF ADMINISTRATION OF THE MEDICINAL PRODUCT:

Treatment should be around 20 days at the average recommended dosage. Between 2 and 6 times a day or as prescribed by the physician.

DURATION OF TREATMENT:

1 Bottle of 5mL allows a treatment of around 20 days at the average recommended dosage. Even after improvement of symptoms, the treatment should be prolonged for the whole period of risk of allergy.

UNWANTED AND TROUBLESOME EFFECTS (UNDESIRABLE EFFECTS):

Possible transient burning or stinging sensations upon instillation.

NOTIFY YOUR DOCTOR OR PHARMACIST OF ANY UNEXPECTED OR TROUBLESOME EFFECTS NOT MENTIONED IN THIS LEAFLET.

STORAGE:

Store in temperatures not exceeding 25°C. This medicinal product should not be kept beyond 4 weeks following the date of opening.

DO NOT EXCEED THE EXPIRY DATE CLEARLY INDICATED ON THE OUTER PACKAGING.

SPECIAL PRECAUTIONS FOR STORAGE:

AVAILABILITY:

5mL HDPE bottle (Box of 1 's)

CAUTION:

Foods, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription.

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph

Manufactured by:

EXCELVISION
27 Rue de la Lombardière,
Annonay, 07100, France

For:

LABORATOIRES
Théa

12 rue Louis Bleriot
63017 Clermont –Ferrand
Cedex 2-France

Imported and Distributed by:

STADA Philippines Inc.
Unit 1001-1003 The Finance Centre
26th Street corner 9th Avenue
Bonifacio Global City
Fort Bonifacio
Taguig City, Metro Manila

Repacked by:

RYTPACK PHARMA ENTERPRISES
G/F Natividad Bldg., 2308 Chino Roces Ave.,
Makati, Metro Manila